

# Activation of invariant natural killer T cells by $\alpha$ -galactosylceramide ameliorates myocardial ischemia/reperfusion injury in mice

*by* Ali Sobirin

---

**Submission date:** 18-Sep-2019 09:00AM (UTC+0700)

**Submission ID:** 1174822463

**File name:** ameliorates\_myocardial\_ischemia\_reperfusion\_injury\_in\_mice..pdf (1.32M)

**Word count:** 8421

**Character count:** 41509



Contents lists available at SciVerse ScienceDirect

## Journal of Molecular and Cellular Cardiology

journal homepage: [www.elsevier.com/locate/yjmcc](http://www.elsevier.com/locate/yjmcc)

## Original article

# Activation of invariant natural killer T cells by $\alpha$ -galactosylceramide ameliorates myocardial ischemia/reperfusion injury in mice



Tsuneaki Homma<sup>a</sup>, Shintaro Kinugawa<sup>a,\*</sup>, Masashige Takahashi<sup>c</sup>, Mochamad Ali Sobirin<sup>a,b</sup>, Akimichi Saito<sup>a</sup>, Arata Fukushima<sup>a</sup>, Tadashi Suga<sup>a</sup>, Shingo Takada<sup>a</sup>, Tomoyasu Kadoguchi<sup>a</sup>, Yoshihiro Masaki<sup>a</sup>, Takaaki Furihata<sup>a</sup>, Masaru Taniguchi<sup>a</sup>, Toshinori Nakayama<sup>d</sup>, Naoki Ishimori<sup>a</sup>, Kazuya Iwabuchi<sup>e</sup>, Hiroyuki Tsutsui<sup>a</sup>

<sup>a</sup> Department of Cardiovascular Medicine, Hokkaido University Graduate School of Medicine, Sapporo, Japan<sup>b</sup> Faculty of Medicine, Diponegoro University, Semarang, Indonesia<sup>c</sup> RIKEN Research Center for Allergy and Immunology, Kanagawa, Japan<sup>d</sup> Department of Immunology, Graduate School of Medicine, Chiba University, Chiba, Japan<sup>e</sup> Department of Immunology, Kitasato University School of Medicine, Sagami-hara, Japan

## ARTICLE INFO

## Article history:

Received 4 February 2013

Received in revised form 21 May 2013

Accepted 5 June 2013

Available online 14 June 2013

## Keywords:

Invariant natural killer T cells

Myocardial ischemia/reperfusion injury

Inflammation

Cytokines

## ABSTRACT

Invariant natural killer T (iNKT) cells orchestrate tissue inflammation via regulating various cytokine productions. However the role of iNKT cells has not been determined in myocardial ischemia/reperfusion (I/R) injury. The purpose of this study was to examine whether the activation of iNKT cells by  $\alpha$ -galactosylceramide ( $\alpha$ -GC), which specifically activates iNKT cells, could affect myocardial I/R injury. I/R or sham operation was performed in male C57BL/6J mice. I/R mice received the injection of either  $\alpha$ -GC (I/R +  $\alpha$ -GC,  $n = 48$ ) or vehicle (I/R + vehicle,  $n = 49$ ) 30 min before reperfusion. After 24 h, infarct size/area at risk was smaller in I/R +  $\alpha$ -GC than in I/R + vehicle ( $37.8 \pm 2.7\%$  vs.  $47.1 \pm 2.5\%$ ,  $P < 0.05$ ), with no significant changes in area at risk. The numbers of infiltrating myeloperoxidase- and CD3-positive cells were lower in I/R +  $\alpha$ -GC. Apoptosis evaluated by TUNEL staining and caspase-3 protein was also attenuated in I/R +  $\alpha$ -GC. Myocardial gene expression of tumor necrosis factor- $\alpha$  and interleukin (IL)-1 $\beta$  in I/R +  $\alpha$ -GC was lower to 46% and 80% of that in I/R + vehicle, respectively, whereas IL-10, IL-4, and interferon (IFN)- $\gamma$  were higher in I/R +  $\alpha$ -GC than I/R + vehicle by 2.0, 4.1, and 9.6 folds, respectively. The administration of anti-IL-10 receptor antibody into I/R +  $\alpha$ -GC abolished the protective effects of  $\alpha$ -GC on I/R injury (infarct size/area at risk:  $53.1 \pm 5.2\%$  vs.  $37.4 \pm 3.5\%$ ,  $P < 0.05$ ). In contrast, anti-IL-4 and anti-IFN- $\gamma$  antibodies did not exert such effects. In conclusion, activated iNKT cells by  $\alpha$ -GC play a protective role against myocardial I/R injury through the enhanced expression of IL-10. Therapies designed to activate iNKT cells might be beneficial to protect the heart from I/R injury.

© 2013 Elsevier Ltd. All rights reserved.

## 1. Introduction

Early and successful myocardial reperfusion is the most effective strategy to reduce infarct size and preserve cardiac function after acute myocardial infarction (MI) [1]. Reperfusion after ischemia can salvage the ischemic myocardium, however, simultaneously it causes

additional cell death and attenuates the beneficial effects of reperfusion itself, called myocardial ischemia/reperfusion (I/R) injury [2]. Inflammation has been shown to play a critical role in the pathophysiology of myocardial I/R injury [3], and various immune cells, such as neutrophils, T lymphocytes, monocytes/macrophages, and mast cells, are involved in myocardial I/R injury [4–7]. Recent study by Yang et al. demonstrated that CD4<sup>+</sup> T lymphocytes played an important role in the development of I/R injury and interferon (IFN)- $\gamma$  was involved in their action by using Rag1 knockout mice lacking mature lymphocytes [5].

Invariant natural killer T (iNKT or type 1 NKT) cells are innate-like T lymphocyte population characterized by co-expressing NK lineage receptors and T cell receptors (TCR), and their TCR has invariant  $\alpha$ -chain (V $\alpha$ 14-J $\alpha$ 18 in mice, and V $\alpha$ 24-J $\alpha$ 18 in humans) [8,9]. They are activated by recognizing glycolipid antigens presented by CD1d, a member of major histocompatibility complex (MHC) class I

**Abbreviations:** AAR, area at risk;  $\alpha$ -GC,  $\alpha$ -galactosylceramide; IFN- $\gamma$ , interferon- $\gamma$ ; IL, interleukin; iNKT, invariant natural killer T; I/R, ischemia/reperfusion; IS, infarct size; LV, left ventricle; MI, myocardial infarction; MNCs, mononuclear cells; MPO, myeloperoxidase; NK, natural killer; qRT-PCR, quantitative reverse transcriptase-polymerase chain reaction; TCR, T cell receptor; T<sub>H</sub>1, T-helper type 1; T<sub>H</sub>2, T-helper type 2; TGF- $\beta$ , transforming growth factor- $\beta$ ; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; TTC, 2,3,5-triphenyltetrazolium chloride.

\* Corresponding author at: Department of Cardiovascular Medicine, Hokkaido University Graduate School of Medicine, Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan. Tel.: +81 11 706 6973; fax: +81 11 706 7874.

E-mail address: [tuckahoe@med.hokudai.ac.jp](mailto:tuckahoe@med.hokudai.ac.jp) (S. Kinugawa).

0022-2828/\$ – see front matter © 2013 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.yjmcc.2013.06.004>



These molecules, and rapidly secrete a mixture of large amount of T-helper type 1 (T<sub>H</sub>1) and T<sub>H</sub>2 cytokines, such as IFN- $\gamma$ , interleukin (IL)-10 and IL-4 in shaping subsequent adaptive immune responses [10]. Thus, iNKT cells can function as a bridge between the innate and adaptive immune systems, and orchestrate tissue inflammation.

iNKT cells have been demonstrated to play a protective role in various autoimmune and inflammatory diseases such as type 1 diabetes, experimental allergic encephalomyelitis, rheumatoid arthritis, and enteritis [11–15]. We have also reported that the activation of iNKT cells by  $\alpha$ -galactosylceramide ( $\alpha$ GC), a specific activator for iNKT cells [16], can attenuate the development of left ventricular (LV) remodeling and failure after MI created by chronic ligation of coronary artery in mice [17]. The activation of iNKT cells by  $\alpha$ GC has also been reported to protect the liver against I/R injury in mice via IL-13 production [18]. However, no previous studies have examined the effects of iNKT cell activation by  $\alpha$ GC on myocardial I/R injury.

Therefore, the purpose of the present study was to determine whether the activation of iNKT cells by  $\alpha$ GC could attenuate myocardial I/R injury. We also determined whether the protective effects on attenuated myocardial I/R injury might involve the activation of anti-inflammatory cytokines including IL-10.

## 2. Materials and methods

Detailed methods are available in the Online Supplementary Material.

### 2.1. Animals

C57BL/6J mice were purchased from CLEA Japan, Inc. (Tokyo, Japan). Animals were used for experiments at 10 to 12 weeks of age (weight 23–27 g). Mice were bred in a pathogen-free environment and kept under a constant 12-h light–dark cycle at a temperature of 23 °C to 25 °C. Standard chow and water were provided.

All procedures and animal care were approved by our institutional animal research committee and conformed to the animal care guideline for the Care and Use of Laboratory Animals in Hokkaido University Graduate School of Medicine.

### 2.2. Experimental design

#### 2.2.1. Experiment 1: effects of $\alpha$ GC on iNKT cell and cytokine expression in the normal mice

To confirm that  $\alpha$ GC could activate iNKT cells in the heart similar to the spleen, C57BL/6J mice were sacrificed 0, 24, and 72 h after  $\alpha$ GC (Funakoshi Co., Ltd., Tokyo, Japan) injection (0.1  $\mu$ g/g body weight i.p.,  $n = 9$  for each group) and the proportion of iNKT cells in the heart and spleen were measured by flow cytometric analysis [17].

To determine that  $\alpha$ GC could induce the changes of cytokines in the blood and the heart within 24 h, another group of C57BL/6J mice were sacrificed 0, 0.5, 1, 3, 6, 12, and 24 h after single injection of  $\alpha$ GC ( $n = 6$  for each group). Serum levels of IL-10, IL-4, and IFN- $\gamma$  were measured by ELISA and their gene expressions in the heart were measured by quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR). TCR in iNKT cells has invariant  $\alpha$ -chain encoded by V $\alpha$ 14-J $\alpha$ 18 gene segment in mice, and J $\alpha$ 18<sup>−/−</sup> mice lack iNKT cells. To confirm whether the changes of cytokines by the injection of  $\alpha$ GC were due to the activation of iNKT cells, iNKT cell-deficient (J $\alpha$ 18<sup>−/−</sup>) mice were sacrificed after single injection of  $\alpha$ GC and same measurements were performed. They were provided from Dr. M. Taniguchi (RIKEN, Yokohama, Japan) and backcrossed 10 times to C57BL/6J.

#### 2.2.2. Experiment 2: effects of $\alpha$ GC on myocardial I/R injury

Myocardial I/R surgery or sham operation was performed in C57BL/6J mice according to the methods described previously [5]. After anesthesia, the left coronary artery was ligated for 45 min.

Ischemia was confirmed by bleaching of the myocardium. Reperfusion was initiated by releasing the ligature. Sham-operated mice underwent a similar procedure without ligation.  $\alpha$ GC (0.1  $\mu$ g/g body weight i.p.) was administered 30 min before reperfusion to specifically activate iNKT cells. As control, the same volume of vehicle was administered into sham and I/R mice.

Mice were sacrificed 24 h after reperfusion (sham + vehicle,  $n = 22$ ; sham +  $\alpha$ GC,  $n = 22$ ; I/R + vehicle,  $n = 49$ ; I/R +  $\alpha$ GC,  $n = 48$ ). These mice were divided into groups for some measurements. Another groups of mice were sacrificed 72 h after reperfusion for flow cytometric analysis ( $n = 9$  for each group), because iNKT cells have been reported to be invisible by flow-cytometric detection 24 h after  $\alpha$ GC administration [19]. Additional mice were sacrificed 72 h after reperfusion for RT-PCR analysis ( $n = 7$ –8 for each group). To confirm early protective effect of  $\alpha$ GC, C57BL/6J mice received I/R surgery with vehicle or  $\alpha$ GC, and sacrificed 2 h after reperfusion to measure infarct size (I/R + vehicle,  $n = 7$ ; I/R +  $\alpha$ GC,  $n = 8$ ). Furthermore, to confirm the effect of  $\alpha$ GC-induced reduction of infarct size on long-term LV function and remodeling, echocardiography and hemodynamic measurement were performed at 28 days after reperfusion (I/R + vehicle,  $n = 8$ ; I/R +  $\alpha$ GC,  $n = 8$ ).

To confirm whether the effect of  $\alpha$ GC on infarct size in I/R was due to the activation of iNKT cells, J $\alpha$ 18<sup>−/−</sup> mice received I/R surgery with vehicle or  $\alpha$ GC and sacrificed 24 h after reperfusion to measure infarct size ( $n = 5$  for each group).

Furthermore, to examine the role of various cytokines in the effects of  $\alpha$ GC on myocardial I/R injury, rat anti-IL-10 receptor monoclonal antibody (200  $\mu$ g/mouse, i.p., BD Pharmingen, San Diego, CA), rat anti-IL-4 monoclonal antibody (250  $\mu$ g/mouse, i.p., R&D System, Inc.), or rat anti-IFN- $\gamma$  monoclonal antibody (150  $\mu$ g/mouse, i.p., R&D System, Inc.) was administered 90 min before I/R surgery and infarct size was measured 24 h after reperfusion. The doses of these antibodies were chosen based on the previous study of their efficacy [18,19,20]. We also confirmed that the changes of serum IL-4 or IFN- $\gamma$  levels were completely inhibited by identical antibodies. Rat IgG1 $\kappa$  was used as control.  $\alpha$ GC was administered 30 min before reperfusion (I/R +  $\alpha$ GC + rat IgG1 $\kappa$ ,  $n = 8$ ; I/R +  $\alpha$ GC + anti-IL-10R,  $n = 8$ ; I/R +  $\alpha$ GC + anti-IL-4,  $n = 7$ ; I/R +  $\alpha$ GC + anti-IFN- $\gamma$ ,  $n = 9$ ).

Finally, to examine the role of IFN- $\gamma$  on myocardial I/R injury, rat anti-IFN- $\gamma$  monoclonal antibody (150  $\mu$ g/mouse, i.p., R&D System, Inc.) or IgG1 $\kappa$  was administered 90 min before I/R surgery and infarct size was measured 24 h after reperfusion ( $n = 6$  for each).

### 2.3. Statistical analysis

Data are expressed as means  $\pm$  SE. The Student  $t$  test was performed for comparison between 2 independent groups. For multiple-group comparisons, one-way ANOVA followed by the Dunnett's test or the Tukey's test was performed. A value of  $P < 0.05$  was considered statistically significant.

The authors had full access to and take full responsibility for the integrity of the data. All authors had read and agree to the manuscript as written.

## 3. Results

### 3.1. Experiment 1: effects of $\alpha$ GC on iNKT cell and cytokine expression in the normal mice

#### 3.1.1. Proportion of iNKT cells after $\alpha$ GC administration

After  $\alpha$ GC administration, splenic iNKT cells disappeared at 24 h and were increased at 72 h (Supplemental Fig. 1, upper panel) in consistency with the previous report [19]. The number of cardiac iNKT cells itself was lower than that of splenic iNKT cells. However, they were increased 72 h after  $\alpha$ GC administration in parallel to splenic

iNKT cells (Supplemental Fig. 1, lower panel). Similar results were observed in 3 independent experiments.

### 3.2. Serum levels and myocardial gene expression of cytokines after $\alpha$ GC administration

After the administration of  $\alpha$ GC, serum IL-10 and IL-4 levels were rapidly increased and peaked at 1 h and 3 h respectively, and serum IFN- $\gamma$  levels were increased later and peaked at 12 h (Supplemental Fig. 2A), which was consistent with the previous reports [21,22]. Gene expression of IL-10, IL-4, and IFN- $\gamma$  in the LV was increased within 24 h and peaked at later phase than serum levels (Supplemental Fig. 2B).

### 3.3. Specificity of $\alpha$ GC for iNKT cells

$\alpha$ GC did not increase serum levels and myocardial gene expression of IL-10, IL-4, and IFN- $\gamma$  in  $J\alpha 18^{-/-}$  mice (Fig. 1).

### 3.4. Experiment 2: effects of $\alpha$ GC on myocardial I/R injury

#### 3.4.1 Body weight and hemodynamics

There was no difference in body weight among all groups. Systolic blood pressure was significantly lower in I/R mice compared to sham mice, however, which was not affected by  $\alpha$ GC. Diastolic blood pressure and heart rate did not differ among 4 groups.

### 3.5. iNKT cells

Representative flow cytometric analyses from 4 groups of mice are shown in Fig. 2A. The proportion of iNKT cells 72 h after reperfusion was increased up to 2.7-fold in I/R + vehicle compared to sham + vehicle.  $\alpha$ GC significantly increased these proportion of iNKT cells both in sham up to 17.5-fold ( $P = 0.042$ ) and I/R mice up to

10.3-fold ( $P = 0.004$ ) (Fig. 2B). Similar results were observed in 3 independent experiments.


### 3.6. I/R injury and LV function

The administration of  $\alpha$ GC into I/R mice decreased infarct size. IS/AAR was significantly smaller in I/R +  $\alpha$ GC than in I/R + vehicle ( $37.8 \pm 2.7\%$  vs.  $47.1 \pm 2.5\%$ ,  $P = 0.018$ ) without significant changes in AAR/LV ( $58.5 \pm 2.3\%$  vs.  $58.9 \pm 2.7\%$ ,  $P = \text{NS}$ ) at 24 h after reperfusion (Fig. 3A). In consistent with these results, serum level of troponin-I was also lower in I/R +  $\alpha$ GC than in I/R + vehicle ( $5.8 \pm 0.8$  ng/mL vs.  $8.7 \pm 0.7$  ng/mL,  $P = 0.016$ , Fig. 3B). At 2 h after reperfusion, IS/AAR was also smaller in I/R +  $\alpha$ GC than in I/R + vehicle ( $20.7 \pm 1.8\%$  vs.  $29.9 \pm 2.8\%$ ,  $P = 0.014$ ) without significant changes in AAR/LV (Supplemental Fig. 3). To examine whether the reduction in infarct size by  $\alpha$ GC was due to the activation of iNKT cells,  $J\alpha 18^{-/-}$  mice were used. There were no differences in IS/AAR between  $J\alpha 18^{-/-}$  + I/R +  $\alpha$ GC and  $J\alpha 18^{-/-}$  + I/R + vehicle ( $34.9 \pm 4.7\%$  vs.  $35.6 \pm 4.1\%$ ,  $P = \text{NS}$ ) and in AAR/LV between groups ( $51.3 \pm 6.4\%$  vs.  $59.0 \pm 2.4\%$ ,  $P = \text{NS}$ ) at 24 h after reperfusion (Supplemental Fig. 4).

Echocardiography and hemodynamic data at 28 days after reperfusion were shown in Table 1. LV end-diastolic dimension did not differ between the 2 groups, whereas LV end-systolic dimension was significantly decreased in I/R +  $\alpha$ GC compared to I/R + vehicle, which resulted in greater fractional shortening in I/R +  $\alpha$ GC. Furthermore, anterior wall thickness including infarct region was preserved in I/R +  $\alpha$ GC. There were no differences in HR, BP, and LV  $\pm$  dP/dt between groups. LV end-diastolic pressure was decreased in I/R +  $\alpha$ GC.


### 3.7. Inflammatory cell infiltration

Immunohistochemical analysis revealed that the number of MPO (as a marker of neutrophil)- and CD3 (as a marker of T lymphocyte)-positive



**Fig. 1.** Specificity of  $\alpha$ GC for iNKT cells. (A) Serum levels of IL-10, IL-4 and IFN- $\gamma$  at 0, 0.5, 1, 3, 6, 12 and 24 h after  $\alpha$ GC intraperitoneal injection into  $J\alpha 18^{-/-}$  mice (0.1  $\mu$ g/g body weight).  $n = 3$  for each group. (B) Quantitative analysis of IL-10, IL-4 and IFN- $\gamma$  mRNA expression in the myocardium after  $\alpha$ GC injection into  $J\alpha 18^{-/-}$  mice.  $n = 3$  for each group. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. 0 h. ND, not detected.






**Fig. 2.** The proportion of iNKT cells in the heart from 4 groups of mice. (A) Representative flow cytometric analyses of cardiac mononuclear cell (MNC) suspensions 72 h after reperfusion from sham + vehicle, sham + αGC, I/R + vehicle and I/R + αGC. Cardiac MNCs from 3 different mice for each group were pooled and analyzed. Circles indicate the population of iNKT cells. (B) Summary data for the proportion of iNKT cells.  $n = 3$  for each group. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. sham + vehicle. † $P < 0.05$  vs. I/R + vehicle.

cells and the ratio of MAC3 (as a marker of macrophage)-positive area in the ischemic myocardium were increased in I/R + vehicle compared to sham + vehicle. Administration of αGC into I/R mice significantly ameliorated the infiltration of MPO- and CD3-positive cells in I/R mice. In contrast, there were no significant differences in MAC3-positive area between I/R + vehicle and I/R + αGC (Figs. 4A, B).

Flow cytometric analysis also showed that CD45<sup>+</sup> cells (leukocytes), CD45<sup>+</sup>Ly6G<sup>+</sup> cells (neutrophils), and Ly6Chigh monocytes in the ischemic heart were decreased in I/R + αGC compared to I/R + vehicle (Supplemental Fig. 5). In contrast, CD45<sup>+</sup>CD68<sup>+</sup> cells excluding monocytes (macrophages) were comparable between groups (Supplemental Fig. 5).

### 3.8. TUNEL staining and caspase-3 protein

There were rare TUNEL-positive nuclei both in sham + vehicle and sham + αGC mice. There were some cardiomyocytes with TUNEL-positive nuclei in I/R + vehicle and I/R + αGC (Fig. 5A). The number of TUNEL-positive cardiomyocytes in the ischemic LV was increased in I/R + vehicle compared to sham + vehicle. It was significantly decreased in I/R + αGC compared to I/R + vehicle (Fig. 5B). Full length caspase-3 protein levels were significantly increased in the ischemic myocardium from I/R + vehicle compared to sham + vehicle, which was consistent with previous papers [23,24], and this increase was also inhibited in I/R + αGC (Fig. 5C).



**Fig. 3.** Effects of αGC on myocardial I/R injury at 24 h. (A) Representative pictures of Evans Blue and TTC-stained LV sections from I/R + vehicle and I/R + αGC (upper panels). IS/AAR and AAR/LV 24 h after reperfusion in I/R + vehicle ( $n = 16$ ) and I/R + αGC ( $n = 14$ ) mice (lower panel). (B) Serum levels of troponin-I 24 h after reperfusion in sham + vehicle ( $n = 8$ ), sham + αGC ( $n = 8$ ), I/R + vehicle ( $n = 16$ ) and I/R + αGC ( $n = 14$ ) mice. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. sham + vehicle. † $P < 0.05$  vs. I/R + vehicle. ND, not detected.

### 3.9. Serum and myocardial cytokines, and chemokines

Very small amounts of serum IL-10 and IFN- $\gamma$  levels were detected and IL-4 was not detected in either sham + vehicle or I/R + vehicle mice. In contrast, αGC extremely increased these cytokine levels in both sham and I/R mice (Fig. 6A). At 24 h after reperfusion, gene expression of IL-10, IL-4, and IFN- $\gamma$  in the ischemic myocardium tended to be increased in I/R + vehicle compared to sham + vehicle, and they were significantly increased in I/R + αGC compared to I/R + vehicle (Fig. 6B). These gene expressions were also measured at 72 h after reperfusion in another set of mice. Gene expression of IL-10 was significantly increased in both I/R + vehicle and I/R + αGC, and tended to be increased by αGC in sham and I/R (Supplemental

Fig. 6A). IL-4 and IFN- $\gamma$  were increased in I/R + αGC compared to I/R + vehicle (Supplemental Figs. 6B, C).

Gene expression of TNF- $\alpha$ , IL-1 $\beta$ , and TGF- $\beta$ 1 was significantly increased in I/R + vehicle compared to sham + vehicle. In contrast to IL-10, IL-4, and IFN- $\gamma$ , TNF- $\alpha$ , and IL-1 $\beta$  were lower in I/R + αGC than I/R + vehicle (Fig. 6B). MCP-1, ICAM-1, and VCAM-1 were increased in I/R + vehicle (Supplemental Fig. 7). MCP-1 was decreased in I/R + αGC compared to I/R + vehicle (Supplemental Fig. 7), which could inhibit infiltration of inflammatory cells (Fig. 4 and Supplemental Fig. 5). In contrast, there was no difference in ICAM-1, and VCAM-1 was rather increased in I/R + αGC compared to I/R + vehicle (Supplemental Fig. 7).

### 3.10. Effects of neutralization of IL-10, IL-4, and IFN- $\gamma$ on αGC-treated I/R mice

Representative pictures showed that the administration of anti-IL-10 receptor monoclonal antibody into I/R + αGC mice increased infarct size compared to I/R + αGC + rat IgG1 $\kappa$ . In contrast, the administration of anti-IL-4 and anti-IFN- $\gamma$  did not affect it (Fig. 7, upper panels). IS/AAR was significantly greater in I/R + αGC + anti-IL-10R than I/R + αGC + rat IgG1 $\kappa$  ( $53.1 \pm 5.2\%$  vs.  $37.4 \pm 3.5\%$ ,  $P = 0.046$ ) with no significant changes in AAR/LV ( $54.3 \pm 2.0\%$  vs.  $54.7 \pm 2.9\%$ ,  $P = \text{NS}$ ) (Fig. 7, lower panel).

### 3.11. Effects of neutralization of IFN- $\gamma$ on I/R mice

Representative pictures showed that the administration of anti-IFN- $\gamma$  monoclonal antibody into I/R mice decreased infarct size compared to I/R + rat IgG1 $\kappa$  (Supplemental Fig. 8, upper panels). IS/AAR was significantly smaller in I/R + anti-IFN- $\gamma$  than I/R + rat IgG1 $\kappa$  ( $38.0 \pm 3.7\%$  vs.  $49.7 \pm 1.9\%$ ,  $P = 0.020$ ) with no significant changes in AAR/LV ( $56.4 \pm 4.1\%$  vs.  $57.8 \pm 1.5\%$ ,  $P = \text{NS}$ ) (Supplemental Fig. 8, lower panel).


## 4. Discussion

The present study demonstrated that the activation of iNKT cells by αGC ameliorated myocardial I/R injury, accompanied by the decreases in inflammatory cell infiltration, apoptosis, and pro-inflammatory cytokines. Furthermore, the neutralization of αGC-induced increase in expression of IL-10 by receptor antibody abolished the protective effects of αGC on I/R injury. This is the first report to provide direct evidence for the protective effects of iNKT cell activation by αGC on myocardial I/R injury.

### 4.1. Activation of iNKT cells by αGC in the heart

αGC has been well known to activate iNKT cells, and they rapidly produce various cytokines such as IL-10, IL-4, and IFN- $\gamma$  [25]. In parallel to these changes, iNKT cell-surface receptors, including TCR and NK1.1, become downregulated, which render iNKT cells invisible by flow-cytometric detection [19,26]. The downregulation of TCR remains until at least 24 h. Then, iNKT cells rapidly proliferate and increase to the peak level 72 h after αGC administration. The activation of iNKT cells by αGC has been observed in various organs, such as spleen, liver, lung, and kidney [18,19,26–28]. We previously demonstrated that the proportion of iNKT cells was increased within the heart 7 days after αGC administration [17]. In the present study, we confirmed that αGC increased cardiac iNKT cells in parallel to splenic iNKT cells (Supplemental Fig. 1) and rapidly enhanced the expression of cytokine genes within the heart (Supplemental Fig. 2B), however, αGC had no effect in iNKT cell-deficient ( $\alpha 18^{-/-}$ ) mice (Fig. 1). Therefore, these findings indicate that αGC can specifically activate iNKT cells, which results in the production of cytokines in the heart.





**Fig. 4.** Effects of αGC on the number of infiltrating inflammatory cells in ischemic myocardium. (A) Representative photomicrographs of LV sections stained with anti-myeloperoxidase (MPO), anti-CD3 and anti-MAC3 24 h after reperfusion. (B) Summary data for the number of MPO-positive cells and CD3-positive cells and the ratio of MAC3-positive area.  $n = 7-8$  for each group. Data are expressed as means  $\pm$  SE. \*  $P < 0.05$  vs. sham + vehicle. †  $P < 0.05$  vs. I/R + vehicle. HPF, high power field.

#### 4.2. Myocardial I/R injury and cytokines

It is well known that various cytokines are involved in myocardial I/R injury [29,30]. These cytokines are produced by several types of cells, such as neutrophils, lymphocytes, macrophages, and endothelial

cells, and play an important role in the pathogenesis of myocardial I/R injury. Endogenous TNF- $\alpha$  and IL-1 play as a mediator of inflammatory reactions, whereas, IL-10 and TGF- $\beta$  have cardioprotective effects on myocardial I/R injury. Previous studies demonstrated that the blocking of pro-inflammatory cytokines or the administration of cardioprotective cytokines reduced infarct size [31–36]. On the other hand, the increases in IL-4 and IFN- $\gamma$  are characteristic of the activation of iNKT cells [9]. IFN- $\gamma$  has been reported to promote myocardial I/R injury [5], and the effect of IL-4 on myocardial I/R injury has not been elucidated. In the present study, αGC administration decreased infarct size (Fig. 3) and infiltrating inflammatory cells (Fig. 4 and Supplemental Fig. 5) in association with the decrease in the expressions of pro-inflammatory cytokines, TNF- $\alpha$  and IL-1 $\beta$  (Fig. 6B). Simultaneously, serum levels and gene expression of IL-10, IL-4, and IFN- $\gamma$  were increased after αGC administration (Figs. 6A, B).

To determine the role of these cytokines in αGC-induced amelioration of myocardial I/R injury, we neutralized IL-10, IL-4, and IFN- $\gamma$  on αGC-treated I/R mice. Anti-IL-10R monoclonal antibody canceled the protective effects of αGC in I/R mice, but not anti-IL-4 and IFN- $\gamma$  monoclonal antibody (Fig. 7), indicating that IL-10 was involved in the protective effects of αGC in myocardial I/R injury. Yang et al. reported that INF- $\gamma$  had deleterious effects on myocardial I/R injury [5]. We also showed that a single treatment with anti-IFN- $\gamma$  monoclonal antibody reduced infarct size after I/R injury without αGC (Supplemental Fig. 8). In our results, the reduction in infarct size by αGC was the same as that by anti-IFN- $\gamma$  monoclonal antibody, and


**Table 1**

Echocardiogram and hemodynamics 28 days after reperfusion.


	I/R + vehicle	I/R + αGC
Echocardiography	$n = 8$	$n = 8$
LVEDD, mm	$3.7 \pm 0.1$	$3.5 \pm 0.1$
LVESD, mm	$3.0 \pm 0.1$	$2.7 \pm 0.1^{\dagger}$
FS, %	$18.9 \pm 0.9$	$24.0 \pm 1.1^{\dagger}$
AWT, mm	$0.64 \pm 0.02$	$0.71 \pm 0.01^{\dagger}$
PWT, mm	$0.80 \pm 0.02$	$0.82 \pm 0.02$
Hemodynamics	$n = 6$	$n = 7$
Heart rate, bpm	$456 \pm 25$	$449 \pm 20$
Systolic BP, mm Hg	$108.2 \pm 6.2$	$108.1 \pm 3.7$
Diastolic BP, mm Hg	$78.8 \pm 4.7$	$76.5 \pm 1.9$
LVEDP, mm Hg	$5.2 \pm 0.4$	$1.9 \pm 0.3^{\dagger}$
LV + dP/dt, mm Hg/s	$10,633 \pm 1824$	$11,841 \pm 1284$
LV - dP/dt, mm Hg/s	$7365 \pm 1600$	$7286 \pm 1105$

LVEDD indicates left ventricular end-diastolic diameter; LVESD, left ventricular end-systolic diameter; FS, fractional shortening; AWT, anterior wall thickness; PWT, posterior wall thickness; BP, blood pressure; LVEDP, left ventricular end-diastolic pressure. Data are expressed as means  $\pm$  SE.

$^{\dagger} P < 0.05$  vs. I/R + vehicle.




**Fig. 5.** Effects of αGC on apoptosis. (A) Representative photomicrographs of TUNEL-stained LV sections. TUNEL-positive nuclei (green), myoglobin (red), and DAPI (blue). Arrows indicate TUNEL-positive cells. (B) Summary data for the number of TUNEL-positive cells ( $n = 7-8$  for each group). (C) Representative immunoblotting analysis and the summary data for caspase-3/GAPDH.  $n = 6$  for each group. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. sham + vehicle. † $P < 0.05$  vs. I/R + vehicle.



**Fig. 6.** Effects of αGC on serum levels and myocardial gene expression of cytokines. (A) Serum levels of IL-10, IL-4 and IFN-γ 24 h after reperfusion. (B) Quantitative analysis of mRNA expression of IL-10, IL-4, IFN-γ, TNF-α, IL-1β, and TGF-β1 in ischemic myocardium 24 h after reperfusion.  $n = 8$  for each group. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. sham + vehicle. † $P < 0.05$  vs. I/R + vehicle. ND, not detected.





**Fig. 7.** Effects of neutralizing antibodies on αGC-induced amelioration of I/R injury. Representative pictures of Evans Blue and TTC-stained LV sections from I/R + αGC + rat IgG1κ, I/R + αGC + anti-IL-10 receptor mAb (IL-10R), I/R + αGC + anti-IL-4 mAb and I/R + αGC + anti-IFN-γ mAb (upper panels). Summary data for IS/AAR and AAR/LV 24 h of reperfusion (lower panel),  $n = 7-9$  for each group. Data are expressed as means  $\pm$  SE. \* $P < 0.05$  vs. I/R + αGC + rat IgG1κ.

there was no additional reduction in infarct size by αGC and anti-IFN-γ monoclonal antibody. These results suggested that IL-10-dependent beneficial effects of αGC may be due to the inhibition of INF-γ-dependent deleterious effects on myocardial I/R injury. Indeed, it has been reported that IL-10 inhibits the expression of IFN-γ-induced genes [37].

We showed that infarct size was smaller in I/R + αGC than I/R + vehicle also at 2 h after reperfusion (Supplemental Fig. 3). Serum IL-10 was rapidly increased and peaked at 1 h after αGC injection, in contrast, gene expression of IL-10 in the LV was not increased at early phase after αGC injection (Supplemental Fig. 2). Therefore, the improvement in infarct size by αGC may be due to the increased serum IL-10, i.e. systemic activation of iNKT cells.

#### 4.3. Protective effects of IL-10 on myocardial I/R injury

IL-10 is well known as a potent anti-inflammatory cytokine [38], and has been shown to play an important role in myocardial I/R injury [33]. It has been reported that endogenous IL-10 inhibits the production of TNF-α and serves to protect the reperfused myocardium through the suppression of neutrophil recruitment [33]. IL-10 has been reported to suppress the expression of CC chemokine gene including MCP-1 [39]. Exogenous IL-10 administration ameliorates myocardial I/R injury by inhibiting adherence of leukocytes to vascular endothelium [34], and by decreasing the production of pro-inflammatory cytokines through Signal Transducers and Activator of Transcription (STAT)-3 pathway [35]. It has also been shown that remote ischemic preconditioning has protective effects against myocardial I/R injury by the upregulation of IL-10 in the remote muscle and the release into circulation [40]. Moreover, IL-10 induces protection against myocardial injury by preventing apoptosis through the reduced phosphorylation of p38MAPK and the enhanced phosphorylation of STAT3 [41]. TNF-α and IL-1β promotes apoptosis in cardiac myocytes [42,43], which is also inhibited by IL-10. In the present

study, we demonstrated that αGC administration ameliorated myocardial I/R injury (Fig. 3) with upregulating serum and myocardial IL-10 (Fig. 6). This was accompanied with the decreases in the infiltration of inflammatory cells into myocardium (Fig. 4 and Supplemental Fig. 5) and the gene expression of pro-inflammatory cytokines (Fig. 6B), and the reduction in apoptosis after I/R (Fig. 5). Our results suggest that the activated iNKT cells by αGC inhibit inflammatory response and cardiomyocyte apoptosis via the production of IL-10.

We previously demonstrated that administration of αGC into mice 1 day and 4 days after MI surgery ameliorated LV remodeling without affecting infarct size and these beneficial effects were also mediated by the enhanced expression of IL-10 in the heart [17]. On the other hand, in the present study, we demonstrated that αGC administration decreased infarct size. The discrepancy in the effects of αGC on infarct size was possibly due to the differences in experimental model assessing different pathophysiological processes (more angiogenesis, more fibrosis vs. accentuated inflammation and apoptosis in the acute setting) and the timing of αGC treatment (1 day after MI surgery vs. 30 min before reperfusion). The administration of αGC into MI mice was performed too late to salvage ischemic myocardium.

#### 4.4. Clinical implication

The present study demonstrated that αGC administration during the ischemic period before reperfusion reduced infarct size. These findings suggest that αGC can be a novel agent in patients with acute MI to reduce I/R injury. In addition, based on our previous study of postinfarct heart failure [17], αGC administration may attenuate also LV remodeling and reduce mortality after MI. To date, several clinical trials (Phase I/II) using activated iNKT cells by αGC have been conducted in patients with cancer [44–48]. No severe adverse events were observed in these trials.



#### 4.5. Limitations

There are several limitations to be acknowledged in the present study. First, we could not directly demonstrate the activation of iNKT cells within the heart 24 h after  $\alpha$ GC administration because cell-surface receptors were downregulated. We tried double immunohistochemical staining of anti-TCR $\beta$  and anti-NK1.1 according to the newly published paper [49]. Furthermore, we also performed in situ hybridization using digoxigenin-labeled DNA probes for mouse V $\alpha$ 14J $\alpha$ 18. Unfortunately, however, we could not detect iNKT cells by these methods in the heart. Further studies are needed to overcome some technical difficulties of iNKT cell-detection and clarify this important issue. Alternatively, we demonstrated its activation by showing the increase of cytokine gene expressions and the similar time course of iNKT cell proportion within the spleen. Second, the source of IL-10 production after the stimulation of  $\alpha$ GC remains to be determined. We tried to isolate iNKT cells using cell sorter and perform in vitro experiments. Unfortunately, however, we could not isolate sufficient amount of iNKT cells to perform in vitro experiments. We consider that there still might be some technical difficulties in in vitro experiments using isolated iNKT cells. Therefore, we could not directly demonstrate the source of IL-10 production. IL-10 has been shown to be produced by iNKT cells themselves on exogenous stimulation [50]. However,  $\alpha$ GC-activated iNKT cells may stimulate other immune cells to produce IL-10. IL-10 can be also expressed and secreted from macrophages activated by iNKT cells. Moreover, in myocardial I/R injury, it has reported that CD5 positive T lymphocytes are the predominant source of IL-10 in the ischemic and reperfused heart. However, immunohistochemical analysis and flow cytometric analysis revealed that there was no difference in the infiltration of macrophage in the ischemic myocardium between I/R +  $\alpha$ GC and I/R + vehicle, and other inflammatory cells were rather decreased in I/R +  $\alpha$ GC compared to I/R + vehicle (Fig. 4 and Supplemental Fig. 5). Further investigations are required to elucidate the mechanism of IL-10 production after  $\alpha$ GC administration in I/R mice. Third, we observed that iNKT cells were increased and endogenously activated in ischemic myocardium after I/R, however, the clear evidence on the role of iNKT cells in I/R injury has not been shown in the present study. We performed I/R injury experiment using iNKT cell deficient J $\alpha$ 18 $^{-/-}$  mice. Unexpectedly, preliminary results showed that infarct size after I/R injury tended to be decreased in J $\alpha$ 18 $^{-/-}$  mice compared to C57BL/6J control mice (IS/AAR in I/R + vehicle group from Fig. 3 vs. IS/AAR in J $\alpha$ 18 $^{-/-}$  + I/R + vehicle group from Supplemental Fig. 4). These results suggest that endogenously activated iNKT cells may be involved in the development of I/R injury, even though endogenous ligand for the activation of iNKT cells has never been elucidated. Therefore, the discrepancy in these results may be due to the difference in the methods to activate iNKT cells; endogenous ligand vs. exogenously administrated  $\alpha$ GC. Finally, protein levels of several cytokines could not be detected in the heart by ELISA in the present study. This may be due to short half-life, and smaller amount of cytokines protein in the heart than in the serum.

#### 5. Conclusions

Activated iNKT cells by  $\alpha$ GC play a protective role against myocardial I/R injury through the enhanced expression of IL-10. Therapies designed to activate iNKT cells might be beneficial to protect the heart from I/R injury.

#### Disclosures

None.

#### Funding

This study was supported by grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (24659379, 20117004, 21390236).

#### Acknowledgments

We thank Akiko Aita, Miwako Fujii and Kaoruko Naradate for excellent technical assistance.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.jmcc.2013.06.004>.

#### References

- [1] Cannon CP, Gibson CM, Lambrew CT, Shultz DA, Levy D, French WJ, et al. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA* 2000;283:2941–7.
- [2] Yellon DM, Hausenloy DJ. Myocardial reperfusion injury. *N Engl J Med* 2007;357:1121–35.
- [3] Zuidema MY, Zhang C. Ischemia/reperfusion injury: the role of immune cells. *World J Cardiol* 2010;2:325–32.
- [4] Hawkins HK, Entman ML, Zhu JY, Youker KA, Berens K, Dore M, et al. Acute inflammatory reaction after myocardial ischemic injury and reperfusion. Development and use of a neutrophil-specific antibody. *Am J Pathol* 1996;148:1957–69.
- [5] Yang Z, Day YJ, Toufektsian MC, Xu Y, Ramos SI, Marshall MA, et al. Myocardial infarct-sparing effect of adenosine A2A receptor activation is due to its action on CD4+ T lymphocytes. *Circulation* 2006;114:2056–64.
- [6] Birdsall HH, Green DM, Trial J, Youker KA, Burns AR, MacKay CR, et al. Complement C5a, TGF-beta 1, and MCP-1, in sequence, induce migration of monocytes into ischemic canine myocardium within the first one to five hours after reperfusion. *Circulation* 1997;95:684–92.
- [7] Frangogiannis NG, Lindsey ML, Michael LH, Youker KA, Bressler RB, Mendoza LH, et al. Resident cardiac mast cells degranulate and release preformed TNF-alpha, initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion. *Circulation* 1998;98:699–710.
- [8] Godfrey DI, MacDonald HR, Kronenberg M, Smyth MJ, Van Kaer L. NKT cells: what's in a name? *Nat Rev Immunol* 2004;4:231–7.
- [9] Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol* 2007;25:297–336.
- [10] Matsuda JL, Mallevaey T, Scott-Browne J, Gapin L. CD1d-restricted iNKT cells, the 'Swiss-Army knife' of the immune system. *Curr Opin Immunol* 2008;20:358–68.
- [11] Hong S, Wilson MT, Serizawa I, Wu L, Singh N, Naidenko OV, et al. The natural killer T-cell ligand alpha-galactosylceramide prevents autoimmune diabetes in non-obese diabetic mice. *Nat Med* 2001;7:1052–6.
- [12] Sharif S, Arreaza GA, Zucker P, Mi QS, Sondhi J, Naidenko OV, et al. Activation of natural killer T cells by alpha-galactosylceramide treatment prevents the onset and recurrence of autoimmune Type 1 diabetes. *Nat Med* 2001;7:1057–62.
- [13] Singh AK, Wilson MT, Hong S, Olivares-Villagomez D, Du C, Stanic AK, et al. Natural killer T cell activation protects mice against experimental autoimmune encephalomyelitis. *J Exp Med* 2001;194:1801–11.
- [14] Miellot A, Zhu R, Diem S, Boissier MC, Herbelin A, Bessis N. Activation of invariant NK T cells protects against experimental rheumatoid arthritis by an IL-10-dependent pathway. *Eur J Immunol* 2005;35:3704–13.
- [15] Ronet C, Darche S, Leite de Moraes M, Miyake S, Yamamura T, Louis JA, et al. NKT cells are critical for the initiation of an inflammatory bowel response against *Toxoplasma gondii*. *J Immunol* 2005;175:899–908.
- [16] Kawano T, Cui J, Koezuka Y, Toura I, Kaneko Y, Motoki K, et al. CD1d-restricted and TCR-mediated activation of valpha14 NKT cells by glycosylceramides. *Science* 1997;278:1626–9.
- [17] Sobirin MA, Kinugawa S, Takahashi M, Fukushima A, Homma T, Ono T, et al. Activation of natural killer T cells ameliorates postinfarct cardiac remodeling and failure in mice. *Circ Res* 2012;111:1037–47.
- [18] Cao Z, Yuan Y, Jeyabalan G, Du Q, Tsung A, Geller DA, et al. Preactivation of NKT cells with alpha-GalCer protects against hepatic ischemia–reperfusion injury in mouse by a mechanism involving IL-13 and adenosine A2A receptor. *Am J Physiol Gastrointest Liver Physiol* 2009;297:G249–58.
- [19] Crowe NY, Uldrich AP, Kyprissoudis K, Hammond KJ, Hayakawa Y, Sidobre S, et al. Glycolipid antigen drives rapid expansion and sustained cytokine production by NK T cells. *J Immunol* 2003;171:4020–7.
- [20] Mabalirajan U, Dinda AK, Kumar S, Roshan R, Gupta P, Sharma SK, et al. Mitochondrial structural changes and dysfunction are associated with experimental allergic asthma. *J Immunol* 2008;181:3540–8.
- [21] Miyamoto K, Miyake S, Yamamura T. A synthetic glycolipid prevents autoimmune encephalomyelitis by inducing TH2 bias of natural killer T cells. *Nature* 2001;413:531–4.



- [22] Parekh VV, Singh AK, Wilson MT, Olivares-Villagomez D, Bezbradica JS, Inazawa H, et al. Quantitative and qualitative differences in the in vivo response of NKT cells to distinct alpha- and beta-anomeric glycolipids. *J Immunol* 2004;173:3693–706.
- [23] Duan W, Yang Y, Yan J, Yu S, Liu J, Zhou J, et al. The effects of curcumin post-treatment against myocardial ischemia and reperfusion by activation of the JAK2/STAT3 signaling pathway. *Basic Res Cardiol*;107:263.
- [24] Bharti S, Golechha M, Kumari S, Siddiqui KM, Arya DS. Akt/GSK-3beta/eNOS phosphorylation arbitrates safranal-induced myocardial protection against ischemia-reperfusion injury in rats. *Eur J Nutr*;51:719–727.
- [25] Van Kaer L.  $\alpha$ -Galactosylceramide therapy for autoimmune diseases: prospects and obstacles. *Nat Rev Immunol* 2005;5:31–42.
- [26] Wilson MT, Johansson C, Olivares-Villagomez D, Singh AK, Stanic AK, Wang CR, et al. The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion. *Proc Natl Acad Sci U S A* 2003;100:10913–8.
- [27] Aoyagi T, Yamamoto N, Hattai M, Tanno D, Miyazato A, Ishii K, et al. Activation of pulmonary invariant NKT cells leads to exacerbation of acute lung injury caused by LPS through local production of IFN-gamma and TNF-alpha by Gr-1 + monocytes. *Int Immunol* 2011;23:97–108.
- [28] Li L, Huang L, Vergis AL, Ye H, Bajwa A, Narayan V, et al. IL-17 produced by neutrophils regulates IFN-gamma-mediated neutrophil migration in mouse kidney ischemia-reperfusion injury. *J Clin Invest* 2010;120:331–42.
- [29] Frangogiannis NG, Youker KA, Rossen RD, Gwechenberger M, Lindsey MH, Mendoza LH, et al. Cytokines and the microcirculation in ischemia and reperfusion. *J Mol Cell Cardiol* 1998;30:2567–76.
- [30] Zhang M, Chen L. Status of cytokines in ischemia reperfusion induced heart injury. *Cardiovasc Hematol Disord Drug Targets* 2008;8:161–72.
- [31] Gurevitch J, Frolkis I, Yuhay S, Lifschitz-Mercer B, Berger E, Paz Y, et al. Anti-tumor necrosis factor-alpha improves myocardial recovery after ischemia and reperfusion. *J Am Coll Cardiol* 1997;30:1554–61.
- [32] Venkatachalam K, Prabhu SD, Reddy VS, Boylston WH, Valente AJ, Chandrasekar B. Neutralization of interleukin-18 ameliorates ischemia/reperfusion-induced myocardial injury. *J Biol Chem* 2009;284:7853–65.
- [33] Yang Z, Zingarelli B, Szabo C. Crucial role of endogenous interleukin-10 production in myocardial ischemia/reperfusion injury. *Circulation* 2000;101:1019–26.
- [34] Hayward R, Nossuli TO, Scalia R, Lefer AM. Cardioprotective effect of interleukin-10 in murine myocardial ischemia-reperfusion. *Eur J Pharmacol* 1997;334:157–63.
- [35] Manukyan MC, Alvernaz CH, Poynter JA, Wang Y, Brewster BD, Weil BR, et al. Interleukin-10 protects the ischemic heart from reperfusion injury via the STAT3 pathway. *Surgery* 2011;150:231–9.
- [36] Lefer AM, Ma XL, Weyrich AS, Scalia R. Mechanism of the cardioprotective effect of transforming growth factor beta 1 in feline myocardial ischemia and reperfusion. *Proc Natl Acad Sci U S A* 1993;90:1018–22.
- [37] Ito S, Ansari P, Sakatsume M, Dickensheets H, Vazquez N, Donnelly RP, et al. Interleukin-10 inhibits expression of both interferon alpha- and interferon gamma-induced genes by suppressing tyrosine phosphorylation of STAT1. *Blood* 1999;93:1456–63.
- [38] Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol* 2001;19:683–765.
- [39] Kopydlowski KM, Salkowski CA, Cody MJ, van Rooijen N, Major J, Hamilton TA, et al. Regulation of macrophage chemokine expression by lipopolysaccharide in vitro and in vivo. *J Immunol* 1999;163:1537–44.
- [40] Cai ZP, Parajuli N, Zheng X, Becker L. Remote ischemic preconditioning confers late protection against myocardial ischemia-reperfusion injury in mice by upregulating interleukin-10. *Basic Res Cardiol* 2012;107:277–107.
- [41] Krishnamurthy P, Rajasingh J, Lambers E, Qin G, Losordo DW, Kishore R. IL-10 inhibits inflammation and attenuates left ventricular remodeling after myocardial infarction via activation of STAT3 and suppression of HuR. *Circ Res* 2009;104:e9–18.
- [42] Krown KA, Page MT, Nguyen C, Zechner D, Gutierrez V, Comstock KL, et al. Tumor necrosis factor alpha-induced apoptosis in cardiac myocytes. Involvement of the sphingolipid signaling cascade in cardiac cell death. *J Clin Invest* 1996;98:2854–65.
- [43] McTiernan CF, Lemster BH, Frye C, Brooks S, Combes A, Feldman AM. Interleukin-1 beta inhibits phospholamban gene expression in cultured cardiomyocytes. *Circ Res* 1997;81:493–503.
- [44] Giaccone G, Punt CJ, Ando Y, Ruijter R, Nishi N, Peters M, et al. A phase I study of the natural killer T-cell ligand alpha-galactosylceramide (KRN7000) in patients with solid tumors. *Clin Cancer Res* 2002;8:3702–9.
- [45] Nieda M, Okai M, Tazbirkova A, Lin H, Yamaura A, Ide K, et al. Therapeutic activation of Valpha24 + Vbeta11 + NKT cells in human subjects results in highly coordinated secondary activation of acquired and innate immunity. *Blood* 2004;103:383–9.
- [46] Ishikawa A, Motohashi S, Ishikawa E, Fuchida H, Higashino K, Otsuji M, et al. A phase I study of alpha-galactosylceramide (KRN7000)-pulsed dendritic cells in patients with advanced and recurrent non-small cell lung cancer. *Clin Cancer Res* 2005;11:1910–7.
- [47] Uchida T, Horiguchi S, Tanaka Y, Yamamoto H, Kunii N, Motohashi S, et al. Phase I study of alpha-galactosylceramide-pulsed antigen presenting cells administration to the nasal submucosa in unresectable or recurrent head and neck cancer. *Cancer Immunol Immunother* 2008;57:337–45.
- [48] Motohashi S, Nagato K, Kunii N, Yamamoto H, Yamasaki K, Okita K, et al. A phase I-II study of alpha-galactosylceramide-pulsed IL-2/GM-CSF-cultured peripheral blood mononuclear cells in patients with advanced and recurrent non-small cell lung cancer. *J Immunol* 2009;182:2492–501.
- [49] Barral P, Sanchez-Nino MD, van Rooijen N, Cerundolo V, Batista FD. The location of splenic NKT cells favours their rapid activation by blood-borne antigen. *EMBO J* 2012;31:2378–90.
- [50] Sonoda KH, Faunce DE, Taniguchi M, Exley M, Balk S, Stein-Streilein J. NK T cell-derived IL-10 is essential for the differentiation of antigen-specific T regulatory cells in systemic tolerance. *J Immunol* 2001;166:42–50.



# Activation of invariant natural killer T cells by $\alpha$ -galactosylceramide ameliorates myocardial ischemia/reperfusion injury in mice

## ORIGINALITY REPORT

6%

SIMILARITY INDEX

8%

INTERNET SOURCES

8%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[fenicia-events.eu](http://fenicia-events.eu)

Internet Source

1%

2

Wataru Mizushima, Hidehisa Takahashi, Masashi Watanabe, Shintaro Kinugawa et al. "The novel heart-specific RING finger protein 207 is involved in energy metabolism in cardiomyocytes", Journal of Molecular and Cellular Cardiology, 2016

Publication

1%

3

Guo, Cai-xia, Xue Jiang, Xiang-jun Zeng, Hong-xia Wang, Hui-hua Li, Feng-he Du, and Bu-xing Chen. "Soluble receptor for advanced glycation end-products protects against ischemia/reperfusion-induced myocardial apoptosis via regulating the ubiquitin proteasome system", Free Radical Biology and Medicine, 2016.

Publication

1%

Li-qun Zhang. "Increased Poly (ADP-

4

ribosyl)ation in Peripheral Leukocytes and the Reperfused Myocardium Tissue of Rats with Ischemia/Reperfusion Injury : Prevention by 3-Aminobenzamide Treatment", Shock, 01/2012

Publication

1%

5

[atvb.ahajournals.org](http://atvb.ahajournals.org)

Internet Source

1%

6

[physoc.onlinelibrary.wiley.com](http://physoc.onlinelibrary.wiley.com)

Internet Source

1%

7

[www.wjgnet.com](http://www.wjgnet.com)

Internet Source

1%

8

[www.science.gov](http://www.science.gov)

Internet Source

1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On